



الحملة الوطنية لمـكافحة فيروس كورونا (COVID-19)

Home isolation Protocol



Home Isolation for Infected individuals of the general public



Infected patients that are Non-Green shield carrier







غير حاصل على التطعيم حاصل على التطعيم **COVID-19 Vaccinated**



Not Vaccinated

- Isolate for a total of 10 days from diagnosis
- No end of isolation swab required

Infected patients that are Green shield carrier



متعافي Recovered



حاصل على التطعيم **COVID-19 Vaccinated**

- Isolate for a total of 7 days from diagnosis
- No end of isolation swab required




HCWs and EWs Home Isolation Protocol






Infected Health Care Workers and Essential Workers (PCR +ve)





Able to verify test:*
Isolate for 7 Days and come
back on D8 to do an Antigen
test at their institution. If
negative resume duty, if
positive return on D11 after
completing the 10 Days
isolation period



Isolate for a total of **10 Days**

HCWs that are returning must be <u>Asymptomatic</u> OR <u>Improving in Symptoms</u> for at least <u>24h</u> prior to resuming duty

*Able to verify test: Able to test in a institute with an infection control unit.

<u>Note:</u> When there is a severe need for essential HCWs, they can return to work immediately or shorten isolation period to 5 days provided they have one negative Ag test and Asymptomatic or improved symptoms for at least 24hrs (as per <u>CDC recommendation</u>) <u>Refer to P.119</u>



completing the 7 Days

isolation period

^{**} Unable to verify test: Unable to test in a institute with an infection control unit.





الحملة الوطنية لمكافحة فيروس كورونا (COVID-19)

CDC recommendation regarding HCWs Quarantine and Isolation

CDC recommendation regarding HCWs Quarantine and Isolation



Work Restrictions for HCP With SARS-CoV-2 Infection and Exposures

"Up to Date" with all recommended COVID-19 vaccine doses is defined in Stay Up to Date with Your Vaccines | CDC

For more details, including recommendations for healthcare personnel who are immunocompromised, have severe to critical illness, or are within 90 days of prior infection, refer to <u>Interim Guidance for Managing Healthcare Personnel with SARS-CoV-2 Infection or Exposure to SARS-CoV-2</u> (conventional standards) and <u>Strategies to Mitigate Healthcare Personnel Staffing Shortages</u> (contingency and crisis standards).

Work Restrictions for HCP With SARS-CoV-2 Infection

Vaccination Status	Conventional	Contingency	Crisis
Up to Date and Not Up to Date	10 days OR 7 days with negative test [†] , if asymptomatic or mild to moderate illness (with improving symptoms)	5 days with/without negative test, if asymptomatic or mild to moderate illness (with improving symptoms)	No work restriction, with prioritization considerations (e.g., types of patients they care for)

Work Restrictions for Asymptomatic HCP with SARS-CoV-2 Exposures

Vaccination Status	Conventional	Contingency	Crisis
Up to Date	No work restrictions, with negative test on days 1 [‡] and 5–7	No work restriction	No work restriction
Not Up to Date	10 days OR 7 days with negative test [†]	No work restriction with negative tests on days 1 [‡] , 2, 3, & 5–7 (if shortage of tests prioritize Day 1 to 2 and 5-7)	No work restrictions (test if possible)

[†]Negative test result within 48 hours before returning to work

[‡]For calculating day of test: 1) for those with infection consider day of symptom onset (or first positive test if asymptomatic) as day 0; 2) for those with exposure consider day of exposure as day 0

